

# Typical self-affine sets with non-empty interior

Zhou Feng

The Chinese University of Hong Kong

Fractal Geometry, Edinburgh, 3–7 July, 2023

Joint work with De-Jun Feng

# Self-affine set

- Let  $T_1, \dots, T_m$  be  $d \times d$  invertible real matrices with  $\|T_j\| < 1/2$  for  $1 \leq j \leq m$ . Write  $\mathbf{T} := (T_1, \dots, T_m)$ .
- For  $\mathbf{a} = (a_1, \dots, a_m) \in \mathbb{R}^{dm}$ , we consider the affine iterated function system,

$$\{f_j^{\mathbf{a}}(x) = T_j x + a_j\}_{j=1}^m.$$


- It is well known that there is a unique non-empty compact set  $K^{\mathbf{a}}$ , called self-affine set, such that

$$K^{\mathbf{a}} = \bigcup_{j=1}^m f_j^{\mathbf{a}}(K^{\mathbf{a}}).$$

- In this talk, we fix the linear part  $\mathbf{T}$  and study  $K^{\mathbf{a}}$  with the translations  $\mathbf{a}$  changing.

# Research target

For example, some self-affine set  $K^{\mathbf{a}}$  looks like



## Goal

To provide some sufficient conditions on  $\mathbf{T}$  such that  $K^{\mathbf{a}}$  has non-empty interior for  $\mathcal{L}^{dm}$ -a.e. (typical)  $\mathbf{a}$ .

# Motivation

In 1988, Falconer introduced a quantity  $\dim_{\text{AFF}} \mathbf{T}$  called **affinity dimension** which only depends on the **linear part**  $\mathbf{T}$ .

## Classical results

- (Falconer, 1988; Solomyak, 1998) For  $\mathcal{L}^{dm}$ -a.e.  $\mathbf{a}$ ,

$$\dim_{\text{H}} K^{\mathbf{a}} = \dim_{\text{B}} K^{\mathbf{a}} = \min \{d, \dim_{\text{AFF}} \mathbf{T}\}.$$

- (Jordan, Pollicott, and Simon, 2007) If  $\dim_{\text{AFF}} \mathbf{T} > d$ , then  $\mathcal{L}^d(K^{\mathbf{a}}) > 0$  for  $\mathcal{L}^{dm}$ -a.e.  $\mathbf{a}$ .

## Question

How about the **interior** of typical  $K^{\mathbf{a}}$ ?

Although this seems a rather fundamental question, it has hardly been studied.

# Main results: general case

To state the result, we define

$$\gamma(\mathbf{T}) = \inf \left\{ \gamma \geq 0 : \sup_{n \geq 1} \sum_{|I|=n} \alpha_d(T_I)^\gamma |\det T_I| \leq 1 \right\}.$$

where  $\alpha_d(T)$  denotes the smallest singular value of a matrix  $T$  and  $T_I = T_{i_1} \cdots T_{i_n}$  for  $I = i_1 \dots i_n \in \{1, \dots, m\}^n$ .

In short,  $\gamma(\mathbf{T})$  is a quantity only depending on  $\mathbf{T}$ .

## Theorem A

If  $\gamma(\mathbf{T}) > d$ , then  $K^{\mathbf{a}}$  has non-empty interior for  $\mathcal{L}^{dm}$ -a.e.  $\mathbf{a}$ .

# The idea for proving Theorem A

- By a classical result (see e.g. (Mattila, 2015)), it suffices to find measures  $\mu^{\mathbf{a}}$  supported on  $K^{\mathbf{a}}$  such that the Fourier transform  $\widehat{\mu^{\mathbf{a}}}$  satisfies

$$\int_{B_\rho} \int_{\mathbb{R}^d} |\widehat{\mu^{\mathbf{a}}}(\xi)|^2 |\xi|^\gamma d\xi d\mathbf{a} < \infty \quad \text{for some } \gamma > d.$$

- By the transversality arguments of Falconer and Solomyak, and some key inequalities, the problem is reduced to finding some measure on  $\{1, \dots, m\}^{\mathbb{N}}$  with enough regularity.
- The condition  $\gamma(\mathbf{T}) > d$  is discovered and provides such a regular measure.

# Main results: commutative case

## Theorem B

Suppose  $T_i T_j = T_j T_i$  for  $1 \leq i, j \leq m$ . If  $\sum_{j=1}^m |\det T_j|^2 > 1$ , then  $K^a$  has non-empty interior for  $\mathcal{L}^{dm}$ -a.e.  $a$ .

For simplicity, we show the idea of the proof for the homogeneous case where

$$\mathbf{T} = (T, \dots, T).$$

- By symbolic expression,

$$K^a = E^a + T E^a,$$

where  $E^a$  is the self-affine set generated by  $\{T^2 x + a_j\}_{j=1}^m$ .

- Then  $m \cdot |\det T|^2 > 1$  implies typically  $\mathcal{L}^d(E^a) > 0$  (Jordan, Pollicott, and Simon, 2007).
- The proof is finished by Steinhaus theorem.

# An open question

By definition,

$$(1) \quad \gamma(\mathbf{T}) > d \iff \exists n \text{ such that } \sum_{|I|=n} \alpha_d(T_I)^d |\det T_I| > 1$$

$$(2) \quad \dim_{\text{AFF}} \mathbf{T} > 2d \iff \sum_{j=1}^m |\det T_j|^2 > 1$$

$$(3) \quad \dim_{\text{AFF}} \mathbf{T} > d \iff \sum_{j=1}^m |\det T_j| > 1.$$

Recall that (1) and (2) are respectively assumed in Theorems A and B. And (3) implies typically  $\mathcal{L}^d(K^a) > 0$ .


Note (1)  $\implies$  (2)  $\implies$  (3).

## Open question


Does typical  $K^a$  have non-empty interior under the condition (3)?




# Some numerical experiments



$\dim_{\text{AFF}} \mathbf{T} < 2$



(3) satisfied



(1) & (2) satisfied

Thank you for listening!



Cheers for fractal!